



**HAL**  
open science

**A comparative analysis of the secondary metabolites, polyphenol contents and antioxidant potential of ethanolic, hydro-ethanolic and aqueous extracts of *Annona muricata* Linn leaves in Abomey-Calavi, Benin**

Isabelle A Appoline, Mahoudo Fidèle Assogba, Lysette Dc Kinsou, Djidénou Ahoton, Mahudjro Yovo, Joachim D Gbenou, Nicolas Atrevi, Patrick A Eдорh

► **To cite this version:**

Isabelle A Appoline, Mahoudo Fidèle Assogba, Lysette Dc Kinsou, Djidénou Ahoton, Mahudjro Yovo, et al.. A comparative analysis of the secondary metabolites, polyphenol contents and antioxidant potential of ethanolic, hydro-ethanolic and aqueous extracts of *Annona muricata* Linn leaves in Abomey-Calavi, Benin. *Journal of Biodiversity and Environmental Sciences* | JBES, 2021, 18 (1), pp.1-10. hal-03578901

**HAL Id: hal-03578901**

**<https://sde.hal.science/hal-03578901>**

Submitted on 17 Feb 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Copyright



## RESEARCH PAPER

## OPEN ACCESS

## A comparative analysis of the secondary metabolites, polyphenol contents and antioxidant potential of ethanolic, hydro-ethanolic and aqueous extracts of *Annona muricata* Linn leaves in Abomey-Calavi, Benin

Appoline Isabelle A Gbonsou<sup>\*1,3</sup>, Mahoudo Fidèle Assogba<sup>2</sup>, Lysette DC Kinsou<sup>2</sup>, Djidénu Ahoton<sup>3</sup>, Mahudjro Yovo<sup>3</sup>, Joachim D Gbenou<sup>1</sup>, Nicolas Atrevi<sup>3</sup>, Patrick A Edorh<sup>1</sup>

<sup>1</sup>Laboratoire de Recherche en Biochimie et Toxicologie de l'Environnement (LaRBiTE), Département de Biochimie et de Biologie Cellulaire, Faculté des Sciences et Techniques (FAST), Université d'Abomey-Calavi (UAC), Cotonou, Bénin

<sup>2,3</sup>Laboratory of Pharmacognosy and Essentials Oils, FAST-FSS, University of Abomey, Calavi, Institute of Applied Biomedical Sciences, Champ de Foire, Cotonou, Benin

<sup>1,3</sup>Laboratoire de Recherche en Biologie Appliquée (LaRBA), Ecole Polytechnique d'Abomey, Calavi (EPAC), Université d'Abomey, Calavi (UAC), Cotonou, Bénin

Article published on January 20, 2021

**Key words:** *Annona muricata*, Leaves, Phytochemical, Antioxidant activity

### Abstract

*Annona muricata* Linn commonly called corossol in BENIN is a plant of the family of Annonaceae known throughout the world through its virtues in traditional medicine. This study aimed to compare the phytochemical composition and the antioxidant activity between ethanolic, hydro-ethanolic, and aqueous extracts of *Annona muricata* leaves collected in Abomey-Calavi, Benin. Qualitative and quantitative phytochemical tests were used to detect the presence of bioactive molecules. Antioxidant activity was performed using the DPPH method and the FRAP reducing power method. The phytochemical screening showed the presence of flavonoids, tannins (catechic and gallic), alkaloid, anthocyanin, leuco anthocyanins, triterpenoids, steroids, mucilage, reducing compounds and coumarins in all three extracts. The ethanolic extract showed a higher antioxidant potential (IC<sub>50</sub>= 0.01mg/ml) than the hydro-ethanolic (IC<sub>50</sub>= 0.013mg/ml) and aqueous extracts (IC<sub>50</sub>= 0.014mg/ml) by the DPPH method. The ethanolic extract showed a higher reducing power than the aqueous extract and the hydro-ethanolic by the FRAP test. This confirms the agreement in the results. However, the ethanolic extract of *Annona muricata* is less active than ascorbic acid, which is a reference antioxidant.

\*Corresponding Author: Appoline Isabelle A Gbonsou ✉ [bellagbons@gmail.com](mailto:bellagbons@gmail.com)

## Introduction

Herbal medicine remains a first-line solution in developing countries and especially for rural populations. So through the ages, man has been able to count on nature to meet basic needs (Lhuillier, 2007). The African continent has a rich heritage of plants with edible and therapeutic virtues. The use of traditional medicine varies from 70% in Benin to 90% in Burundi and Ethiopia (WHO, 2013).

According to the World Health Organization (WHO), medicinal plants are a good source for the discovery and development of new drug of new medicines. Therefore, medicinal plants should be studied to better understand their different properties, safety and efficacy (Nascimento *et al.*, 2000). In Benin, of the 3,000 plant species inventoried in forest ecosystems, 172 are consumed by local populations as food plants and 814 as medicinal plants (Sinsin B., Owolabi L., 2001).

Among these plants that constitute the wealth of the Beninese forest is the corossol tree. Commonly referred to by the name of its fruit, the corossol "magic fruit". It is said to originate in tropical America and then spread to other tropical regions of the world: Europe, Asia, Africa (Gavamukulya *et al.*, 2014; Pieme *et al.*, 2014). The soursop tree with the scientific name *Annona muricata* Linn belongs to the large family of Annonaceae. All the organs of the soursop tree (pulp, stem, leaf, bark, root, seed) are used in herbal medicine (Padma *et al.*, 1999).

Numerous scientific researches carried out in the plant have validated pharmacological activities including hepatoprotective (Padma *et al.*, 1999), hypotensive (Nwokocha *et al.*, 2012), anti-inflammatory (Abdul Wahab *et al.*, 2018), insomnia (Yajid *et al.*, 2018), antiviral, hypoglycemic and antidiabetic (Adewole et Caxton-Martins, 2009; Florence *et al.*, 2014), antioxydant (George *et al.*, 2015), insecticide (Leatemia et Isman, 2004), (Ishuwa *et al.*, 2016). The soursop tree also possesses bioactive substances, antioxidants and rich in acetogenins that give it a potential biological activity and even an anti-cancerous property

(Ana *et al.*, 2016; Coria-Télez *et al.*, 2018; Widyastuti et Rahayu, 2017).

The leaves are used to prevent and overcome arthritis, asthma, bronchitis, biliary disorders, diabetes, heart disease, hypertension, intestinal worms, liver disorders, malaria, rheumatism, tumours and cancers (Gavamukulya *et al.*, 2014; Pieme *et al.*, 2014), (Orak *et al.*, 2019).

Considering the therapeutic importance given to this plant, the present study aimed to make a comparative analysis of the phytochemical composition and the potential antiradical power of the ethanolic, hydroethanolic and aqueous extracts of the leaves of *Annona muricata* Linn harvested in Benin.

## Material and methods

### 1.1. Vegetal material

The vegetal material consists of the leaves of *Annona muricata* Linn, harvested in October 2019 in the district of Toba in the commune of Abomey-Calavi. They have been certified at the National Herbarium of Benin under the number YH264/HNB. The leaves were dried at laboratory temperature (25°C-30°C) for 45 days. Once dried, they were powdered.

### Preparation of extracts

The technique used is that of maceration. 50g of sample *Annona muricata* leaves powder were introduced into a 1000ml bottle containing 500ml of the extraction solvents (Ethanol, water and 50/50 ethanol-water). The vial was capped and left under continuous shaking for 72 hours. After filtration, the extracts were evaporated dry at 60°C using a Heidolph type rotavapor. The yield (R) of the extraction is calculated by the formula below.

$$R (\%) = \frac{\text{Mass of the extract}}{\text{Mass of plant product used}} \times 100$$

### 1.3. Determination of Secondary Metabolites

Phytochemical screening is based on the differential reactions (colouring and precipitation) of the main groups of chemical compounds contained in plants according to the method of (Houghton et Raman, 1998), taken up by (Assogba, *et al.*, 2015).

### *Determination of phenolic compounds*

#### *Total Phenols*

The determination of total polyphenols was carried out according to the method of Singleton and Rossi, 1965, taken up by Ahoton *et al.*, 2019 with some slight modifications. This assay is carried out using the Folin-Ciocalteu reagent and the content is expressed in milligrams of gallic acid equivalent per gram of extract.

#### *Flavonoids*

The total flavonoid content of plant extracts can be estimated by the aluminium trichloride ( $\text{AlCl}_3$ ) method. Rutin is used as the reference compound to produce the calibration curve (Kim DO *et al.*, 2003). The content is expressed in milligram of Rutin equivalent per gram of extract.

#### *Tannins*

The method used to determine the content of condensed tannins is that of sulphuric vanillin (Broadhurst et Jones, 1978 and modified by (Heimler *et al.*, 2006). The condensed tannin content is expressed in milligrams of pyrogallol equivalent per gram of extract.

#### *Anti- Radical Activity*

The antiradical activity of the extracts was determined using two methods: the DPPH free radical scavenging method and the FRAP iron reduction method.

#### *DPPH free radical scavenging test*

For this test, samples were prepared by dissolution in distilled water (Ahoton & al., 2019). For each extract, a stock solution was prepared in distilled water at  $200\mu\text{g/mL}$ . This solution is then diluted in a geometric series of 2 parts to different concentrations.

In dry, sterile test tubes, 1ml of the test extract solution is introduced, 1ml of DPPH solution ( $0.04\text{mg/ml}$ ) is added. After vortex shaking, the tubes are placed in a dark place at laboratory temperature for 30 min. The absorbance measurement is performed at  $517\text{nm}$  with the spectrophotometer (Biomate UV/VIS).

For each dilution, a blank is prepared consisting of 1ml of the test solution plus 1ml of ethanol. The positive control is represented by ascorbic acid ( $200\mu\text{g/ml}$ ) and is treated under the same conditions as the test sample.

The results were expressed as the mean of three measurements  $\pm$  standard deviation. The values of  $\text{IC}_{50}$  (the concentration of the substrate that causes 50% loss of DPPH activity),  $\text{EC}_{50}$  (the effective concentration of the substrate that causes 50% loss of DPPH activity and takes into account the concentration of DPPH in the reaction medium) and APR (antiradical potency) were determined (Parakash D *et al.*, 2007).

#### *Iron Reduction Test (FRAP)*

The extract ability to reduce  $\text{Fe}^{3+}$  was assessed (Oyaizu, 1986). 1ml of the extract was mixed with 2.5ml phosphate buffer (0.2 M, pH 6.6) and 2.5ml of 1% potassium ferricyanide [ $\text{K}_3\text{Fe}(\text{CN})_6$ ]. After a 20 min incubation at  $50^\circ\text{C}$ , 2.5ml of trichloroacetic acid (10%) were added to the mixture and centrifuged for 10 min (3000 r/t). A 2.5ml aliquot of the upper layer was mixed with 2.5ml of distilled water and  $0.5\text{FeCl}_3$  (0.1%). The absorbance was read at  $700\text{nm}$ . Iron (III) reducing activity was determined as ascorbic acid equivalents (mmol ascorbic acid/g extract). The values are presented as a mean of triplicate analyses.

#### *Statistical Analysis*

The data were entered and processed in Excel and Graph Pad 5 software for comparison of means, determination of the confidence index and Pearson correlation coefficients.

### **Results discussion**

#### *Extract yield*

The Table 1 present the yield values for the different extracts.

The hydro-ethanol and aqueous extracts gave the highest yields; the ethanolic extract gave the lowest yield. Yield values differed from each other with a statically significant difference ( $p < 0.05$ ). In Brazil, high percentages were found (12.5%) for ethanol extract and for water extract (15%) (Justino *et al.*, 2018).

Our percentages, although low, are higher than the yields obtained by Orak *et al* 2019 in Turkey, which were solvent dependent (for hexane (3.66%) dichloromethane (1,10%) and ethyl acetate (0.83%)) but lower than for methanolic extraction (12.07%) (Orak *et al.*, 2019). Ethanolic extraction obtained by Ojezele *et al.* (2016) in Nigeria gave a yield (4.9%) lower than our result (Ojezele *et al.*, 2016). Nga *et al.* (2018) in Cameroon obtained high yields for hydro-ethanolic and aqueous extractions respectively (18.07% and 11.87%) (Nga *et al.*, 2018), The quality of the solvent, the harvesting region of the organ used, the drying time, the granulometry of the grind, the percentage of ethanol in the hydro-ethanolic solvent, the ratio of solvent volume per mass of grind, the maceration time and the stirring speed are all factors that influence the yield (Koné *et al.*, 2017).

**Table 1.** Extraction yields.

Extracts	Mass of plant product used (g)	Mass of extract (g)	Yields (%)
Ethanolic	50	3,854	7,708
Hydro-ethanolic	50	4,418	8,836
Aqueous	50	4,162	8,324

### 3.2. Phytochemical screening

The phytochemical screening results of *Annona muricata* extracts are put in the table 2. The table 2 presents bioactive compounds such as alkaloids, flavonoids, reducing compounds, coumarins which are present in all three extracts, soponosides, anthracenes, and cardenolides are absent in all three extracts. The flavonoids identified are flavones. The same compounds are found in the ethanol extracts of the leaves of *Annona muricata* in Nigeria and Andonesia (Ojezele *et al.*, 2016), (Hasmila *et al.*, 2019). The methanolic extraction performed in Nigeria revealed the presence of the cardioglycoside absent in our study (Ezealisiji et Belema, 2017). Cardioglycosides were also found in hydroalcoholic and aqueous leaves extracts in Cameroon (Nga *et al.*, 2018). Other authors have highlighted these different metabolites in different organs of *Annona muricata* (Vijayameena *et al.*, 2013), (Ana *et al.*, 2016), (Onyegeme-Okerenta *et al.*, 2018). However, differences in the absence or presence of any of these

metabolites in different studies could be explained by the solvent used, the identification methodology, the age of the plant, the nature of the soil, the climatic conditions in the country, the region of harvest within the same country. In vivo and in vitro studies have demonstrated the pharmacological activity of the leaves and other organs of the plant due to the presence of flavonoids, tannins, alkaloids and its richness in acetogenin. It has been shown to have antibacterial, antiviral, antifungal, hepatoprotective, anxiolytic and hypoglycemic activity (Florence *et al.*, 2014), (George *et al.*, 2015),(Saleem, 2017),(Ana *et al.*, 2016), (Hansra *et al.*, 2014),(Iyanda-Joel *et al.*, 2019). The anti-tumour properties revealed in breast and prostate tumours were linked to flavonoids and acetogenin richness in the leaves of *Annona muricata* (Gavamukulya *et al.*, 2014; Hansra *et al.*, 2014; Coria-Téllez *et al.*, 2018). These different bioactivities justify the popularity of using this plant in phytotherapy.

**Table 2.** Phytochemical composition of *Annona muricata* Linn.

Chemical compounds	Ethanolic	Hydro-ethanolic	Aqueous extract
Alkaloids	+	+	+
Tannins	+	+	+
Cathechic tannins	-	+	+
Gallic tannins	+	+	+
Flavonoïds	+	+	+
	(flavone)	(flavone)	(flavone)
Anthocyanins	+	-	+
Leucoanthocyan	+	+	-
Quinone derivatives	-	+	+
Mucilage	-	+	+
Reducing compound	+	+	+
Saponins	-	-	-
Coumarins	+	+	+
Free anthracene	-	-	-
O-Heterosides	-	-	-
C-Heterosides	-	-	-
Triterpenoids	+	+	-
Steroids	+	+	-
Cardiac glycosides	-	-	-

(+) Presence; (-) Absence

### 3.3. Determination of phenolic compounds

Polyphenolic compounds dosed in leaf extracts act as antioxidants by inhibiting radicals induced by oxidative stress (Ezealisiji et Belema, 2017). Considering flavonoids, for example, the presence of

the flavone nucleus, the number, positions and types of substitutions influence radical scavenging and chelation activity (Heim *et al.*, 2002). In our work we were able to quantify three chemical families including flavonoids, polyphenols and tannins (fig. 2, 3, 4). Flavonoids in the ethanolic extract are about twice as high as in the other extracts, (fig. 3). but this value is lower than that obtained in Nigeria (365.6 ( $\mu\text{g}/\text{mg}$  QE) with methanolic extracts of *Annona muricata* leaves (Ezealisiji et Belema, 2017). As for polyphenols, they are 1.5 times more expressed in the hydroethanolic extract than in the aqueous and

ethanolic extracts. The same remark was made in Cameroon, with the values of  $22.52 \pm 0.0011$  and  $16.94 \pm 0.0015$  ( $\mu\text{g}/\text{mg}$  GAE) respectively for the hydroethanolic and aqueous it should be noted that polyphenols are higher in our extracts than those of Cameroon (Nga *et al.*, 2018). The maceration time was 72h in our study versus 48 h in Cameroon. The aqueous extract presents relatively low contents compared to the other extracts (fig. 2). In India, the polyphenol content for a concentration of  $100\mu\text{g}$  was  $10.3 \pm 0.06\mu\text{g}/\text{mg}$  GAE (George *et al.*, 2015) value lower than  $75.142 \pm 0.917\mu\text{g}/\text{ml}$  observed in our study.



Fig. 1. Photography of the *Annona muricata* Linn tree with its fruit on the left and dried leaves powder on the right.




Fig. 2. Calibration curve for gallic acid and total phenol contents.






Fig. 3. Rutin calibration curve and flavonoid contents.

*Antiradical activity*

For this activity, we determined the concentration at which half of the DDPH (IC<sub>50</sub>) was reduced by each extract. These IC<sub>50</sub> values are shown in Table 3.

Table 3. DDPH and FRAP tests results.

	IC <sub>50</sub> (µg/ml)	EC <sub>50</sub>	ARP	FRAP (mmol ascorbic acid/g extract)
Aqueous extract	14.01	0.7	1.429	0.182
Hydro-ethanol extract	13.02	0.65	1.538	0.301
Ethanolic extract	10.01	0.5	2	1.126
Ascorbic acid	0.0074	3.7 e-3	8.378	/

Table 3 above shows the values of the concentrations that inhibited 50% of the free radicals in DPPH. These values are indicative of the ability of leaves extracts to scavenge free radicals.

The antioxidant activity of the three extracts was evaluated by two methods DPPH and FRAP for confirmatory purposes. The lower IC<sub>50</sub>, show the more active extract (Heimler *et al.*, 2006). It thus appears from Table 3 that the ethanolic extract is more active than the hydroethanolic extract, which is also more active than the aqueous extract with a statistically significant difference (P<0.05) between the values.

Studies by Coria-Téllez *et al* (2016) in Mexico, by Orak *et al.* (2019) in Turkey, Hasmila *et al.* (2019) in Indonesia showed that ethanolic extract is more active respectively (IC<sub>50</sub> = 70µg/ml and 0.136 ± 0.06mg/ml, 141.127µg/ml). However our extract is the most active.

The ethanolic and aqueous extracts of our study gave lower IC<sub>50</sub> values than those found in Brazil respectively: 28.1±4.4µg/ml and 43.1±6.3µg/ml (Justino *et al.*, 2018), Similarly in Nigeria, the antioxidant activity sought in the ethanolic and aqueous extracts of *Annona muricata* showed that the ethanolic extract is more active 37.8mg/ml versus 79.95mg/ml. Our results corroborate the latter and indicate that the ethanolic extract is more active. However, it is less active than ascorbic acid (0.00749µg/ml).

The iron reduction test (FRAP) is a simple, rapid and reproducible test (Benzie et Strain, 1996). It is universal and can be applied to plants as well as organic and aqueous plasmas and extracts (Ahoton *et al.*, 2019). The reducing power of the extracts has shown that the ethanolic extract of *Annona muricata* has a power three times higher than the hydroethanolic extract and at least five times higher than the aqueous extract which confirms our results with DPPH.


The reducing power of the *Annona muricata* species is probably due to the presence of hydroxyl groups in phenolic compounds that can serve as electron donors. Previous studies have reported that the reducing power of a compound can be a significant indicator of its antioxidant activity (Jeong *et al.*, 2004). The ethanolic extract of the leaves showed a higher antioxidant capacity than the extract of seeds (Widyastuti et Rahayu, 2017).

It was proven that the ethanolic extract has the capacity to decrease the risk of chronic diseases caused by free radicals, in particular cancer and heart attacks also the capacity to reduce the peroxidation of the lipids of the liver (Justino *et al.*, 2018). This radical scavenging capacity could be due to the high flavonoid content in the various extracts. Comparing antiradical activities with total phenol, flavonoid, and tannin levels in the linear correlation model (Pearson test) (Table 4), non-significant correlation was found, with an estimated coefficient of determination, ( $r^2$ ) (fig. 5), at  $p < 0.05$ .


**Table 4.** Correlation between antioxidant activity and content.

	Flavonoïd	Polyphenol	Tannin
IC <sub>50</sub> (mg/mL)	0,7570773	0,3293843	-0,2418611
p-value	0,1214614	0,3353078	0,3790694

The flavonoids presenting the best correlation would be at the base of the antioxidant activity of the extracts. Other authors have shown that phenolic compounds such as flavonoids in plants are the main bioactive substances responsible for antioxidant activity (J *et al.*, 2003), (Ahmed *et al.*, 2015).



**Fig. 4.** Calibration curve for pyrogallol and tannin contents.



**Fig. 5.** Correlation between IC<sub>50</sub> and three chemical groups.

**Conclusion**

Phytochemical screening of the three ethanolic, hydro-ethanolic, and aqueous extracts of *Annona muricata* Linn leaves have increased their wealth in secondary metabolites and phenolic compounds. The Pearson's correlation test allowed us to observe that flavonoids could be at the base of the antioxidant

activity of the extracts. The ethanolic extract has a strong potential to trap free radicals even though it is low in total polyphenols. It is probably for this richness in secondary metabolites that traditional therapists advise the leaves and other organs of the plant in the fight against diseases, especially chronic degenerative diseases.



## References

- Abdul Wahab SM, Jantan I, Haque Md A, Arshad L.** 2018. Exploring the Leaves of *Annona muricata* L. as a Source of Potential Anti-inflammatory and Anticancer Agents. *Frontiers in Pharmacology* **9**. <https://doi.org/10.3389/fphar.2018.01633>
- Adewole S, Caxton-Martins E.** 2009. Morphological changes and hypoglycemic effects of *Annona muricata* Linn. (Annonaceae) leaf aqueous extract on pancreatic  $\beta$ -cells of streptozotocin-treated diabetic rats. *African Journal of Biomedical Research* **9(3)**. <https://doi.org/10.4314/ajbr.v9i3.48903>
- Ahmed D, Khan MM, Saeed R.** 2015. Comparative Analysis of Phenolics, Flavonoids, and Antioxidant and Antibacterial Potential of Methanolic, Hexanic and Aqueous Extracts from *Adiantum caudatum* Leaves. *Antioxidants* **4(2)**, 394. <https://doi.org/10.3390/antiox4020394>
- Ahoton D, Assogba MF, Ladekan ECY, Davo A, Moudachirou M, Gbenou JD.** 2019. Phenolic compounds' dosage and antioxidant activities of aqueous and ethanolic extracts of *Lophira lanceolata*'s Van Tiegh. Ex Keay (Ochnaceae) leaves. *International Journal of Phytopharmacy*.
- Ana C, Montalvo-González E, Yahia E, Obledo-Vázquez E.** 2016. *Annona muricata*: A comprehensive review on its traditional medicinal uses, phytochemicals, pharmacological activities, mechanisms of action and toxicity. *Arabian Journal of Chemistry* **11**. <https://doi.org/10.1016/j.arabjc.2016.01.011>
- Assogba Fidèle M, Aderomou chouaibou, Agbodjogbe Wilfrid, Moudachirou Mansourou, Joachim D Gbenou.** 2015. Evaluation of diuretic properties from *Elaeis guineensis* Jacq. (Arecaceae) leaves aqueous extract in wistar rat [Journal of Chemical and Pharmaceutical research **7(3)**, 2457-2462,]. <https://bec.uac.bj/publication/5631/Article>
- Benzie IFF, Strain JJ.** 1996. The Ferric Reducing Ability of Plasma (FRAP) as a Measure of "Antioxidant Power": The FRAP Assay. *Analytical Biochemistry* **239(1)**, 70-76. <https://doi.org/10.1006/abio.1996.0158>
- Coria-Téllez AV, Montalvo-González E, Yahia EM, Obledo-Vázquez EN.** 2018. *Annona muricata*: A comprehensive review on its traditional medicinal uses, phytochemicals, pharmacological activities, mechanisms of action and toxicity. *Arabian Journal of Chemistry* **11(5)**, 662-691. <https://doi.org/10.1016/j.arabjc.2016.01.004>
- Ezealisiji KM, Belema TE.** 2017. Comparative evaluation of the phenolic and antioxidant properties of the leaves, root, stem bark, and root bark of *Annona muricata* (Annonaceae). *Journal of Pharmacognosy and Phytochemistry* **6(2)**, 274-278.
- Florence NT, Benoit MZ, Jonas K, Alexandra T, Désiré DDP, Pierre K, Théophile D.** 2014. Antidiabetic and antioxidant effects of *Annona muricata* (Annonaceae), aqueous extract on streptozotocin-induced diabetic rats. *Journal of Ethnopharmacology* **151(2)**, 784-790. <https://doi.org/10.1016/j.jep.2013.09.021>
- Gavamukulya Y, Abou-Elella F, Wamunyokoli F, AEl-Shemy H.** 2014. Phytochemical screening, anti-oxidant activity and *in vitro* anticancer potential of ethanolic and water leaves extracts of *Annona muricata* (Graviola). *Asian Pacific Journal of Tropical Medicine* **7S1**, S355-363.
- George VC, Kumar DRN, Suresh PK, Kumar RA.** 2015. Antioxidant, DNA protective efficacy and HPLC analysis of *Annona muricata* (soursop) extracts. *Journal of Food Science and Technology* **52(4)**, 2328-2335.
- Hansra D, Silva O, Mehta A, Ahn E.** 2014. Patient with Metastatic Breast Cancer Achieves Stable Disease for 5 Years on Graviola and Xeloda after Progressing on Multiple Lines of Therapy. *Advances in Breast Cancer Research* **03**, 84-87.
- Hasmila I, Natsir H, Soekamto NH.** 2019. Phytochemical analysis and antioxidant activity of soursop leaf extract (*Annona muricata* Linn.). *Journal of Physics: Conference Series* **1341**, 032027.

- Heimler D, Vignolini P, Dini MG, Vincieri FF, Romani A.** 2006. Antiradical activity and polyphenol composition of local Brassicaceae edible varieties. Food Chemistry. <https://agris.fao.org/agris-search/search.do?recordID=US201300738619>
- Houghton P, Raman A.** 1998. Laboratory handbook for the fractionation of natural extracts. <https://www.google.fr/search?source=hp&ei=BhrSXpyDO6eMlwTGubbQAQ&q=Houghton+P>
- Ishuwa M, Elkanah O, Wahedi J.** 2016. Grain Protectant Potential of *Annona muricata* and *A. senegalensis* against Cowpea Seed Bruchid *Callosobruchus maculatus* Fab. (Coleoptera: Bruchidae). Asian Research Journal of Agriculture **1(1)**, 1-7. <https://doi.org/10.9734/ARJA/2016/2605>
- Iyanda-Joel WO, Ajetunmobi OB, Chinedu SN, Iweala EEJ, Adegbite OS.** 2019. Phytochemical, Antioxidant and Mitochondrial Permeability Transition Studies on Fruit-Skin Ethanol Extract of *Annona muricata* [Research Article]. Journal of Toxicology; Hindawi. <https://doi.org/10.1155/2019>
- JJ, CS, EL, Jm V.** 2003. Antioxidant activity and total phenolic content of Iranian *Ocimum accessions*. Food Chemistry **83(4)**, 547-550. [https://doi.org/10.1016/S0308-8146\(03\)00151-1](https://doi.org/10.1016/S0308-8146(03)00151-1)
- Jeong SM, Kim SY, Kim DR, Jo SC, Nam KC, Ahn DU, Lee SC.** 2004. Effect of heat treatment on the antioxidant activity of extracts from citrus peels. Journal of Agricultural and Food Chemistry **52(11)**, 3389-3393. <https://doi.org/10.1021/jf049899k>
- Justino AB, Miranda NC, Franco RR, Martins MM, Silva NM da, Espindola FS.** 2018. *Annona muricata* Linn. Leaf as a source of antioxidant compounds with in vitro antidiabetic and inhibitory potential against  $\alpha$ -amylase,  $\alpha$ -glucosidase, lipase, non-enzymatic glycation and lipid peroxidation. Biomedicine & Pharmacotherapy **100**, 83-92. <https://doi.org/10.1016/j.biopha.2018.01.172>
- Kim DO Ok C, Yj K, Hy M, Cy L.** 2003. Quantification of polyphenolics and their antioxidant capacity in fresh plums. Journal of Agricultural and Food Chemistry **51(22)**, 6509-6515. <https://doi.org/10.1021/jf0343074>
- Koné K, Yaya S, Sorho S.** 2017. Détermination des paramètres influençant le rendement d'extraction hydro-alcoolique des métabolites secondaires de *Alchornea cordifolia* (Euphorbiaceae) et *Tridax procumbens* Linn (Asteraceae) Détermination des paramètres influençant le rendement d'extraction hydro-alcoolique des métabolites secondaires de *Alchornea cordifolia* (Euphorbiaceae) et *Tridax procumbens* Linn (Asteraceae).
- Leatemia JA, Isman MB.** 2004. Insecticidal activity of crude seed extracts of *Annona* spp., *Lansium domesticum* and *Sandoricum koetjape* against lepidopteran larvae. Phytoparasitica **32(1)**, 30-37.
- Lhuillier A.** 2007. Contribution à l'étude phytochimique de quatre plantes malgaches : *Agauria salicifolia* Hook.f ex Oliver, *Agauria polyphylla* Baker (Ericaceae), *Tambourissa trichophylla* Baker (Monimiaceae) et *Embelia concinna* Baker (Myrsinaceae). [Phd].
- Nascimento GGF, Locatelli J, Freitas PC, Silva GL.** 2000. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. Brazilian Journal of Microbiology **31(4)**, 247-256.
- Nga EN, Nchinda G, Mahi LL, Figuei PB, Mvondo JM, Sagnia B, Adiogo D.** 2018. Phytochemical Screening and Evaluation of Antioxidant Power of Hydro-Ethanollic and Aqueous Leaves Extracts of *Annona muricata* Linn (Soursop). Health Sciences And Disease, **20(1)**, Article 1.
- Nwokocha CR, Owu DU, Gordon A, Thaxter K, McCalla G, Ozolua RI, Young L.** 2012. Possible mechanisms of action of the hypotensive effect of *Annona muricata* (soursop) in normotensive Sprague-Dawley rats. Pharmaceutical Biology **50(11)**, 1436-1441.

- Ojezele O, Ojezele M, Adeosun A.** 2016. Comparative Phytochemistry and Antioxidant Activities of Water and Ethanol Extract of *Annona muricata* Linn Leaf, Seed and Fruit. *Advances in Biological Research* **10**, 230-235. <https://doi.org/10.5829/idosi.abr.2016.10.4.10514>
- Onyegeme-Okerenta BMA, Amadi B, O Ezeonyilimba V.** 2018. The Ameliorating Potential of *Annona muricata* on Sodium Fluoride-induced Toxicity on Liver and Kidney of Male Wistar Rats. *Journal of Complementary and Alternative Medical Research* **6(1)**, 1-17. <https://doi.org/10.9734/JOCAR>
- Orak HH, Bahrisefit IS, Sabudak T.** 2019. Antioxidant Activity of Extracts of Soursop (*Annona muricata* L.) Leaves, Fruit Pulps, Peels, and Seeds. *Polish Journal of Food and Nutrition Sciences* **69(4)**, 359-366. <https://doi.org/10.31883/pjfn/112654>
- Oyaizu M.** 1986. Studies on products of browning reaction—Antioxidative activities of products of browning reaction prepared from glucosamine. *Eiyogaku zasshi= Japanese journal of nutrition*. <https://agris.fao.org/agris-search/search.do?recordID=US201302009163>
- Padma P, Chansouria JPN, Khosa RL.** 1999. Hepatoprotective activity of *Annona muricata* Linn. And *Polyalthia cerasoides* bedd. *Ancient Science of Life* **19(1-2)**, 7-10.
- Parakash D, Brahma UG, Singh N, Singh HB.** 2007. Antioxidant and free radical scavenging activities of seeds and agri-wastes of some varieties of soybean (*Glycine max*) **104**, 783-790.
- Pieme CA, Kumar SG, Dongmo MS, Moukette BM, Boyoum FF, Ngogang JY, Saxena AK.** 2014. Antiproliferative activity and induction of apoptosis by *Annona muricata* (Annonaceae) extract on human cancer cells. *BMC Complementary and Alternative Medicine* **14**. <https://doi.org/10.1186/1472-6882-14-516>
- Saleem Dr U** 2017. Pharmacological Screening of *Annona muricata*: A Review. *Asian Journal of agriculture and biology* **5**, 38-46.
- Sinsin B, Owolabi L.** 2001. Monographie nationale de la diversité biologique. (p. 41) [Rapport de synthèse. MEHU/PNUD].
- Vijayameena C, Subhashini G, Loganayagi M, Balasubramanian R.** 2013. Phytochemical screening and assessment of antibacterial activity for the bioactive compounds in *Annona muricata*. *International Journal for Current Microbiology and Applied Sciences* **2**, 1-8.
- WHO.** 2013. Enhancing the Role of Traditional Medicine in Health Systems: A Strategy for the African Region (p. 13). WHO: Regional Committee For Africa.
- Widyastuti D, Rahayu P.** 2017. Antioxidant Capacity Comparison of Ethanolic Extract of Soursop (*Annona muricata* Linn.) Leaves and Seeds as Cancer Prevention Candidate. *Biology, Medicine, & Natural Product Chemistry* **6**, 1. <https://doi.org/10.14421/biomedich.2017.61.1-4>
- Yajid A, Rahman H, Wong M, Wan Zain W.** 2018. Potential Benefits of *Annona muricata* in Combating Cancer: A Review. *Malaysian Journal of Medical Sciences* **25**, 5-15. <https://doi.org/10.21315/mjms2018.25.1.2>