



A review on potential role of auxins in plants, current applications and future directions

Muhammad Jamil, Amna Saher, Sidra Javed, Qasim Farooq, Muhammad Shakir, Tooba Zafar, Lobaba Komal, Kabir Hussain, Amara Shabir, Arooj Javed, et al.

► To cite this version:

Muhammad Jamil, Amna Saher, Sidra Javed, Qasim Farooq, Muhammad Shakir, et al.. A review on potential role of auxins in plants, current applications and future directions. Journal of Biodiversity and Environmental Sciences | JBES, 2021, 18 (1), pp.11-16. hal-03601670

HAL Id: hal-03601670

<https://sde.hal.science/hal-03601670>

Submitted on 8 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



RESEARCH PAPER

OPEN ACCESS

A review on potential role of auxins in plants, current applications and future directions

Muhammad Jamil¹, Amna Saher², Sidra Javed^{*4}, Qasim Farooq¹, Muhammad Shakir¹, Tooba Zafar¹, Lobaba Komal¹, Kabir Hussain³, Amara Shabir¹, Arooj Javed¹, Muhammad Huzafa¹

¹Department of Botany, University of Agriculture, Faisalabad, Pakistan

²Department of Botany, University of the Punjab, Lahore, Pakistan

³Botany Division, Institute of Pure and Applied Biology, Bahauddin Zakariya University, Multan, Pakistan

⁴Department of Plant Pathology, University of Agriculture, Faisalabad, Pakistan

Article published on January 20, 2021

Key words: Auxin, Plant growth, Apical meristem, Stem elongation, Indole acetic acid

Abstract

Auxin is a plant hormone that significantly plays important role in plant growth cell division and differentiation, in fruit development, in the formation of roots from cuttings and in leaf fall. Auxin has many important functions for plant growth. The main function of auxin is to help the plants grow. Auxin stimulates plant cells to elongate, and the apical meristem of a plant is one of the main places that auxin is produced. It promotes the stem elongation, inhibit growth of lateral buds maintains apical dominance. Auxin is produced in different parts of plants such as the stem, buds, and root tips in the form of chemical compounds such as indole acetic acid that significantly increases the growth of plants. Auxin as a plant hormone produced in the stem tip that promotes cell elongation. Auxin moves to the darker side of the plant, causing the cells there to grow larger than corresponding cells on the lighter side of the plant. Auxin biosynthesis plays essential roles in many developmental processes including gametogenesis, embryogenesis, seedling growth, vascular patterning, and flower development. The aims of this review is the potential roles of auxins in plants such as roles in root development, functions in the growth of leaves structures and lastly the formation of flowers by increases auxins have been studied in this review. The functions of auxins in plants have not completely studied.

***Corresponding Author:** Sidra Javed ✉ sidrajaved943@gmail.com

Introduction

Use of compounds relevant to auxins source on top of the stigmas of tomato and numerous other species causes the ovary to mature into the natural or artificially induced production of fruit without fertilization of ovules (Mapelli *et al.*, 1978). The use of pollen extracts to the external part of the ovary indicated same results, which led to the theory that pollen grains comprise plant hormones similar to the plant growth substance auxin. After fertilization, the pollen may pass enough amount of these plant growth substances to the ovary to initiate growth of fruit (Wang *et al.*, 2018).

Now subsequently, gibberellic acid like plant growth substances were recognized in diverse flowering plants families, thus assuming that these plant growth substances are also involved in the programme of fruit development. It was recognized that concentrations of auxin in the flowers with a traditional bioassay were cut and put on agar blocks. Consequently, the content of auxin in these blocks was evaluated by the standard *Avena* geo-curvature test. No auxin was present at the stage of anthesis. On the other hand, concentration of auxin improved within twenty eight after gibberellic acid treatment, proposing that it is not auxin, but gibberellic acid that is passed on from the germinating pollen to the ovary. Consequently, the gibberellic acid may induce a rise of the content of auxin in the ovary to stages sufficient to activate growth of fruit (Jong *et al.*, 2009).

In spermatophytes, the principal body axes of plants are patterned both during all the life cycle and early in development of embryo. The embryonic development causes the axis of apical-basal, the radial axis, the embryonic leaf and the meristems of primary plumule and radicle. During period of development after embryogenesis, these meristems produce plant lateral organs along the primary growth and create the progress zone. By comparing with animals, plants are thus able to develop morphemes during the whole lifespan. These modular morphological elements can also differ in form in reaction to adaptable environmental signals. Phenotypic plasticity in plant

growth influence consequently restrain evolution in a very diverse way from animals inasmuch as the ultimate shape of the entire plant is not so channeled. In flowering plants, all these chiefmodelling events involve the auxin containing plant growth substances. Furthermore, auxin facilitates growth of plant in reaction to environmental signals. Therefore, the evolution of auxin homeostasis and response systems is supposed to show a main part in the evolution of land plant structural design (Cooke *et al.*, 2004).

The aims of this review is the potential roles of auxins in plants such as roles in root development, functions in the growth of leaves structures, response of auxins in biotic stress in plants, role of auxins transcriptional level in plants, the formation of flowers by increases auxins have been discussed in this review. The functions of auxins in plants have not completely studied.




Fig. 1. Various functions of auxins for plant growth.

Auxin is Essential for the Flowers Formation

Auxin has long been assumed as an important controller for development of flower, but the precise auxin role in development of flower was not well clear in recent times. A connection between auxin and development of flower was first recognized when the mutant of auxin transport *pin1* was isolated and categorized (Gallavotti *et al.*, 2013).

The mutant *pin1* showed pleiotropic deficiencies during the course of practically all phases of development, however the most outstanding phenotype is that the inflorescence of *pin1* regularly does not have any

flowers proposing that auxin shows a vital part in originating flower primordia (Li *et al.*, 2019).

Our understanding of how auxin biosynthesis and degradation are regulated in the root system under normal and different stress conditions is still very limited, but it is clear that auxin homeostasis has a major impact on root architecture. Many genes believed to be involved in auxin biosynthesis are strongly expressed in the root apical meristem, where there is a high rate of IAA biosynthesis. Furthermore, local auxin biosynthesis combined with polar auxin transport is involved in forming the auxin gradient within the root apex. Auxin signaling is key to many plant growth and developmental processes from embryogenesis to senescence. Most, if not all, of these processes are initiated and/or mediated through auxin-regulated gene expression. Pin1 translates a protein of transmembrane that has recently been revealed to localize with polarity in cells and to contribute in auxin efflux (Cheng *et al.*, 2007).




Fig. 2. Role of Auxins in the formation of flowers.

Role of Indole-3-acetic acid in Floral Organ Patterning

Indole-3-acetic acid not only regulates whether flower primordia are formed, it also shows a crucial part in identifying floral organs and is responsible for the developmental process within a floral organ. Numerous *Arabidopsis* mutants were initially isolated from mutagenesis screen for mutants with abnormal flowers, but later found to be involved in aspects of indole-3-acetic acid pathways (Pfluger *et al.*, 2004). Wild-type of flowers of *Arabidopsis* generally have six stamens, four sepals, two fused carpels and four petals.

Alongside the orientation of apical/basal axis, wild-type female part of flower can be categorized into three sections with well-known structures: a basal ovary, a style and an apical stigma (Damodharan *et al.*, 2016).

The allelic alterations affect the growth of all four kinds of floral organs, but the maximum intense deficiency is in female part of flower patterning. Alterations in the gene *ETTIN* intensely affect female part of flower with decrease in the size of ovary and improper position of style of flower and stigma (Van *et al.*, 2012).




Fig. 3. Role of Auxins in the formation different parts of flower.

The chief driver of body plan multiformity is signified by the activity and formation leaf primordia of axillary buds, processes that as defined above are extremely linked to biology of auxin. Biosynthesis, signaling and transport of auxin are all essential for initiation of axillary meristem, as established by diverse defective mutants in these pathways in numerous species, i.e. *mp*, *yuc* and *pin1* in *Arabidopsis*, and *bif2*, *spi* in *Zea mays*, which fail to form new tissues (Den Herder *et al.*, 2010).

While leaf primordia of axillary buds outgrowth has greater consideration in the previous few years with the findings of the complex interaction among cytokinins, auxin and strigolactones quiet relatively little is well-known about how auxin stimulates the

formation of new leaf primordia of axillary buds, although the huge number of categorized mutants that are affected in their initiation (Tsunoda and van Dam, 2017). Leaf primordia of axillary buds initiation can thus work as a model example to study how pathways of auxin interrelated among themselves and with other pathway of developmental process (Waite *et al.*, 2020).

Role of auxin-responsive genes in responding to biotic stress

At molecular level to recognize the potential role of auxin in plant defense responses, the auxin-responsive genes expression was examined under numerous biotic stress circumstances in *Oryza sativa*. Patterns of differential auxin accumulation within a field of cells determine spatial and temporal patterns of auxin-dependent developmental reprogramming. Nonetheless, in the end, the decision on the type of developmental output lies in the interpretation of these auxin accumulations at the level of the individual cell. Despite the diversity of cellular responses evoked by auxin, most of the effects of auxin come down to one relatively simple pathway that directly impinges on transcriptional regulation. In essence, it is the interplay between two classes of transcriptional regulators that represents the core of auxin signaling (Xu *et al.*, 2018).

The DNA microarray data for *Oryza sativa* seedlings infected with *Magnaporthe grisea*, a sac fungi, and *Striga hermonthica*, an obligate root hemiparasite, available at gene expression data and hybridization arrays, chips, microarrays database under series variety numbers GSE10373 and GSE7256, individually, was used for this examination. Under GSE7256 series based on microarray data for two-week old seedlings of *Oryza sativa* variety Nipponbare cultivar treated with haemacytometer of *Magnaporthe grisea* genes for pathogenicity and virulence strain FR13 in gelatin alone and contains 8 hybridizations signifying two biologically distinct samples of infected and mock samples for two time phases (3 and 4 dpi) (Emenecker *et al.*, 2020).

The series GSE10373 based on microarray data of root tissues from three-week old seedlings of two *Oryza sativa* varieties IAC165 (susceptible) and Nipponbare cultivar (resistant) treated with *Striga hermonthica* and contains of twenty-four hybridizations representing two biologically distinct samples of infected and mock samples for three time phases (Meng *et al.*, 2019).




Fig. 4. Transcriptional Roles of auxins in plants.

Roles of auxins in different biotic interactions affecting the root systems

Plant–microorganism associations have gotten increasingly more consideration because of the advantages they present to edit efficiency by improving supplement take-up, expanding plant development and giving biotic and abiotic stress.

Recognizing correspondence frameworks and signs that decide the valuable or impeding results of plant–microorganism collaborations is a key to improve guard reactions without diminishing gainful (e.g., harmonious) affiliations (Du and Scheres *et al.*, 2018).

Auxin influx carriers stabilize phyllotactic patterning

One of the most prominent characteristics of architecture of plant is the symmetrical alignment of leaves and flowers around the stem. Peaks in concentration of the indole-3-acetic acid, created by the polar localization of the efflux of PIN1 auxin

carrier, gives an instructive signal for initiation of primordium. This tool makes the spacing among adjacent primordia, that results in symmetrical alignment of leaves and flowers around the stem (Lee *et al.*, 2015).




Fig. 5. Roles of auxins in different biotic interactions affecting the root systems.

Analysis of the role of indole-3-acetic acid transport in phyllo tactic arrangement have focused on efflux of PIN1 auxin carrier. Current computer models specify an additional role for indole-3-acetic acid transport uptake. Alterations in the influx of AUX1 auxin carrier have not, on the other hand, been described to cause an aerial phenotype. At this time, we studied the role of AUX1 and its gene copies LAX1, LAX2, and LAX3. Study of the four mutant tells asymmetrical deviation angles among succeeding primordia. An extremely rare feature of the phenotype is the existence of groups of primordia, in violation of traditional theory (Bainbridge *et al.*, 2008).

At the molecular level, the well-defined sharp peaks in level of auxin and corresponding PIN polarization are lost. In adding, the augmented penetrance of the phenotype under short-day circumstances recommends that the AUX LAX transporters act to buffer the PIN-mediated model mechanism in contrast to environmental or developmental influences (Julien *et al.*, 2019).




Fig. 6. Roles of auxins influx carriers stabilize phyllotactic patterning.

Conclusion

Several key auxin biosynthetic genes and their roles in plant development have been discovered. It is clear that auxin biosynthesis takes place locally in response to both environmental and developmental signals. *De novo* auxin biosynthesis plays an essential role in many developmental processes. The power of auxin in plants is clearly wide and deep. To understand it, we need to find the missing component parts of the auxin machinery. During the last ten years, our understanding of auxin metabolism and its role during plant growth and development has greatly improved. The great challenge will be to integrate knowledge about auxin metabolism into the regulatory networks that act on different developmental processes operating in plants, and to understand how these processes work in different plant species under normal and stress conditions.

References

- Bainbridge KS, Kuhlemeier C.** 2008. Auxin influx carriers stabilize phyllotactic patterning. *Genes development* **22**, 810-823.
- Cheng Y, Zhao Y.** 2007. A role for auxin in flower development. *Journal of integrative plant biology* **49**, 99-104.

- Cooke TJ, Poli D, Cohen JD.** 2004. Did auxin play a crucial role in the evolution of novel body plans during the Late Silurian-Early Devonian radiation of land plants?, in *The evolution of plant physiology*. Elsevier **55**, 85-107.
- Den GG.** 2010. The roots of a new green revolution. *Trends in plant scienc* **15**, 600-607.
- Emenecker RJ, Strader LC.** 2020. Auxin-Absciscic Acid Interactions in Plant Growth and Development. *Biomolecules* **10**, 281.
- Gallavotti A.** 2013. The role of auxin in shaping shoot architecture. *Journal of experimental botany* **64**, 2593-2608.
- Jong MC, Mariani BG.** 2009. The role of auxin and gibberellin in tomato fruit set. *Journal of experimental botany* **60**, 1523-1532.
- Julien JD, Pumir A, Boudaoud A.** 2019. Strain- or stress-sensing in mechanochemical patterning by the phytohormone auxin. *Bulletin of mathematical biology* **81**, 3342-3361.
- Lee HW, Kim J.** 2015. Lateral Organ Boundaries Domain16 and 18 act downstream of the AUXIN1 and LIKE-AUXIN3 auxin influx carriers to control lateral root development in Arabidopsis. *Plant physiology* **168**, 1792-1806.
- Li T.** 2019. Calcium signals are necessary to establish auxin transporter polarity in a plant stem cell niche. *Nature communications* **10**, 1-9.
- Mapelli S.** 1978. Relationship between set, development and activities of growth regulators in tomato fruits. *Plant and Cell Physiology* **19**, 1281-1288.
- Meng F, Xiang D, Zhu J.** 2019. Molecular mechanisms of root development in rice. *Rice* **12**, 1-10.
- Pflugger J, Zambryski P.** 2004. The role of SEUSS in auxin response and floral organ patterning. *Development* **131**, 4697-4707.
- Wang Y, Jiao Y.** 2018. Auxin and above-ground meristems. *Journal of Experimental Botany* **69**, 147-154.
- Waite JM.** 2020. AtDRO1 is nuclear localized in root tips under native conditions and impacts auxin localization. *Plant molecular biology* **55**, 1-14.
- Xu L.** 2018. De novo root regeneration from leaf explants: wounding, auxin, and cell fate transition. *Current opinion in plant biology* **41**, 39-45.
- Van S.** 2012. Simulation of organ patterning on the floral meristem using a polar auxin transport model. *PloS one* **7**, 155-159.